(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 27 January 2005 (27.01.2005)

(10) International Publication Number WO 2005/007887 A1

(51) International Patent Classification7:

C12Q 1/68

(21) International Application Number:

PCT/GB2004/003086

15 July 2004 (15.07.2004) (22) International Filing Date:

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0316555.2 0328425.4

15 July 2003 (15.07.2003)

8 December 2003 (08.12.2003) GB

(71) Applicant and

(72) Inventor: DENSHAM, Daniel, Henry [GB/GB]; Mobious Genomics Ltd, Innovation Centre, University of Exeter, Rennes Drive, Exeter EX4 4RN (GB).

(74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GII, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MEASUREMENT OF A POLYNUCLEOTIDE AMPLIFICATION REACTION

(57) Abstract: A quantitative measurement of the progress of a polynucleotide amplification reaction can be made by: (i) carrying out a reaction for the amplification of a target polynucleotide; (ii) either during or after the amplification reaction contacting the amplified product with a molecule that binds to or interacts with a polynucleotide, the molecule being located in a spatially defined position or being determined via a non-linear or non-fluorescent technique; and (iii) detecting the interaction between the amplified product and the molecule by measuring changes in radiation.